

Stokes Matrix for the Quantum Cohomology of Cubic Surfaces

Kazushi Ueda

Abstract

We prove the conjectural relation between the Stokes matrix for the quantum cohomology of X and an exceptional collection generating $D^b\text{coh}(X)$ when X is a smooth cubic surface. The proof is based on a toric degeneration of a cubic surface, the Givental's mirror theorem for toric manifolds, and the Picard-Lefschetz theory.

1 Introduction

Let X be a smooth cubic surface in $\mathbb{P}^3$. It is a rational surface obtained by blowing up $\mathbb{P}^2$ at six points. Let

$$\phi : X \rightarrow \mathbb{P}^2$$

denote this blow-up. For a given $\mathbf{d} \in H_2(X; \mathbb{Z})$, let

$$n(\mathbf{d}) = \int_{\mathbf{d}} c_1(TX) - 1$$

be the expected dimension of the moduli space of stable maps of genus zero and degree $\mathbf{d}$ and

$$N_{\mathbf{d}} = \underbrace{\langle [pt], \dots, [pt] \rangle}_{n(\mathbf{d}) \text{ times}}, \mathbf{d}$$

be the Gromov-Witten invariant, where $[pt]$ denotes the Poincaré dual of the homology class of a point. When $n(\mathbf{d}) > 0$, $N_{\mathbf{d}}$ is the number of nodal rational curves in X of degree $\mathbf{d}$ passing through $n(\mathbf{d})$ points in general position. Although it is possible to count $N_{\mathbf{d}}$ by elementary method when the degree $\mathbf{d}$ is small, the difficulty grows rapidly as $\mathbf{d}$ increases.

The theory of Gromov-Witten invariants gives a new perspective to this classical problem of counting $N_{\mathbf{d}}$ in enumerative geometry. Instead of considering $N_{\mathbf{d}}$'s individually, one can treat them as a whole by the following generating function on the cohomology ring $H^*(X; \mathbb{C})$:

$$\Phi(t_0, \dots, t_8) = \frac{1}{2}t_0^2t_8 + t_0(t_1^2 - t_2^2 - \dots - t_7^2) + \sum_{\mathbf{d} \in H_2(X; \mathbb{Z})} N_{\mathbf{d}} q^{\mathbf{d}} \frac{t_8^{n(\mathbf{d})}}{n(\mathbf{d})!}. \quad (1)$$

Here, $t_0, t_1, t_2, \dots, t_7, t_8$ are coordinates on the total cohomology ring $H^*(X; \mathbb{C})$ of X corresponding to the basis $[X], [H], [E_1], \dots, [E_6], [pt]$ of $H^*(X; \mathbb{C})$, where $[X]$ denotes the dual of the fundamental class of X , $[H]$ that of the hyperplane class, and $[E_i]$ that of the exceptional divisor for $i = 1, \dots, 6$, and $q^{\mathbf{d}} = \exp[\langle \mathbf{d}, [H] \rangle t_1] \exp[\langle \mathbf{d}, [E_1] \rangle t_2] \cdots \exp[\langle \mathbf{d}, [E_6] \rangle t_7]$. Then, this Φ endows $H^*(X; \mathbb{C})$ with the structure of a *Frobenius manifold*.

A Frobenius manifold is a complex manifold whose holomorphic tangent bundle has a bilinear form and an associative commutative product structure satisfying a series of axioms. In the case of a Frobenius manifold coming from the Gromov-Witten invariants of a smooth projective variety, the bilinear form is given by the Poincaré pairing, and the structure constant of the product structure is given by the third derivative of the generating function Φ . This product structure is called the quantum cohomology ring, whose associativity gives a highly non-trivial non-linear differential equation for Φ , called the *WDVV (Witten-Dijkgraaf-Verlinde-Verlinde)* equation, which sometimes determines all the genus-zero Gromov-Witten invariants from a finite numbers of them [22].

An important point in the theory of Frobenius manifolds is its connection with isomonodromic deformations. Given a Frobenius manifold, one can construct an isomonodromic family of ordinary differential equations on $\mathbb{P}^1$ of the following form:

$$\hbar \frac{\partial Y}{\partial \hbar} + \frac{1}{\hbar} \mathcal{U}(Y) - \mathcal{V}(Y) = 0, \quad (2)$$

$$\hbar \frac{\partial Y}{\partial t_a} = \frac{\partial}{\partial t_a} \circ Y, \quad a = 0, \dots, N-1. \quad (3)$$

Here, Y is an unknown function on $\mathbb{P}^1$ times the Frobenius manifold taking value in the tangent bundle of the Frobenius manifold, $\hbar$ is the coordinate on $\mathbb{P}^1$, N is the dimension of the Frobenius manifold, $\{t_a\}_{a=0}^{N-1}$ is the *flat coordinate* of the Frobenius manifold, $\circ$ denotes the product on the tangent bundle, and $\mathcal{U}, \mathcal{V}$ are certain operators acting on sections of the tangent bundle. (2) is an ordinary differential equation on $\mathbb{P}^1$ with a regular singularity at infinity

and an irregular singularity at the origin, and (3) gives its isomonodromic deformation. If a point on the Frobenius manifold is semisimple, i.e., if there are no nilpotent elements in the product structure on the tangent space at this point, one can define the *monodromy data* of (2) at this point, consisting of the monodromy matrix at infinity, the Stokes matrix at the origin, and the connection matrix between infinity and the origin. These data do not depend on the choice of a semisimple point because of the isomonodromicity.

The following conjecture, originally due to Kontsevich, developed by Zaslowsky [28], and formulated into the following form by Dubrovin [10], reveals a striking connection between the Gromov-Witten invariants and the derived category of coherent sheaves:

Conjecture 1. *The quantum cohomology of a smooth projective variety X is semisimple if and only if the bounded derived category $D^b\text{coh}(X)$ of coherent sheaves on X is generated as a triangulated category by an exceptional collection $(\mathcal{E}_i)_{i=1}^N$. In such a case, the Stokes matrix S for the quantum cohomology of X is given by*

$$S_{ij} = \sum_k (-1)^k \dim \text{Ext}^k(\mathcal{E}_i, \mathcal{E}_j). \quad (4)$$

An exceptional collection appearing above is the following:

Definition 2. 1. An object $\mathcal{E}$ in a triangulated category is exceptional if

$$\text{Ext}^i(\mathcal{E}, \mathcal{E}) = \begin{cases} \mathbb{C} & \text{if } i = 0, \\ 0 & \text{otherwise.} \end{cases}$$

2. An ordered set of objects $(\mathcal{E}_i)_{i=1}^N$ in a triangulated category is an exceptional collection if each $\mathcal{E}_i$ is exceptional and $\text{Ext}^k(\mathcal{E}_i, \mathcal{E}_j) = 0$ for any $i > j$ and any k .

Conjecture 1 was previously known to hold for projective spaces [11], [17] and Grassmannians [27]. The main result in this paper is:

Theorem 3. *Conjecture 1 holds for smooth cubic surfaces in $\mathbb{P}^3$.*

To prove Theorem 3, we first use the deformation invariance of the Gromov-Witten invariants to reduce the problem to the case of a toric surface Y defined in Section 6 obtained by bringing some of the blowing-up centers to infinitely-near points. Then we can use the Givental's mirror theorem [15], which gives an integral representation of the fundamental solution to (2), (3) for Y of the following form:

$$I_\Gamma = \int_\Gamma \phi(x, y; \hbar) \exp[W(x, y)/\hbar] \frac{dx dy}{xy} \quad (5)$$

Here, $\phi(x, y; \hbar)$ is a certain $H^*(Y; \mathbb{C})$ -valued function of $x, y, \hbar$ and the flat coordinate $\{t_a\}_{a=0}^8$, and $W(x, y)$ is a Laurent polynomial of x and y whose coefficients depend on the flat coordinate. Γ runs over certain cycles in $(\mathbb{C}^\times)^2$ depending on $\hbar$ and flat coordinate, which are descending Morse cycles for $\Re[W(x, y)/\hbar]$ for a suitable choice of a metric on $(\mathbb{C}^\times)^2$.

Since the integrand in (5) is single-valued, the monodromy of the integral comes solely from the monodromy of the integration cycle Γ , which is determined by intersection numbers of vanishing cycles by the Picard-Lefschetz theory. This gives the left-hand side of (4). On the right-hand side, one can find an exceptional collection generating the derived category of coherent sheaves on Y by combining the theorems of Beilinson [4] and Kapranov-Vasserot [21]. The Ext-groups between them can be computed explicitly, and exhibit a complete agreement with the result on the left-hand side.

Acknowledgements: We thank H. Iritani for patiently explaining mirror symmetry for toric manifolds. We also thank A. Ishii, T. Kawai, H. Kawanoue, H. Nasu, K. Saito, A. Takahashi, and H. Uehara for valuable discussions and comments. The author is supported by JSPS Fellowships for Young Scientists No.15-5561.

2 Frobenius manifold and Stokes matrix

We review the relation between Frobenius manifolds and isomonodromic deformations in this section. See, e.g., [9], [11], [18], [19], [25] for more details.

Definition 4. A Frobenius manifold is a quintuple $(M, \circ, e, E, g)$ satisfying the following axioms:

1. M is a complex manifold.
2. $g : \mathcal{T}_M \otimes \mathcal{T}_M \rightarrow \mathcal{O}_M$ is an $\mathcal{O}_M$ -bilinear form.
3. $\circ : \mathcal{T}_M \otimes \mathcal{T}_M \rightarrow \mathcal{T}_M$ defines an associative commutative $\mathcal{O}_M$ -algebra structure on $\mathcal{T}_M$.
4. $g(X \circ Y, Z) = g(X, Y \circ Z)$ as an $\mathcal{O}_M$ -linear map $\mathcal{T}_M \otimes \mathcal{T}_M \otimes \mathcal{T}_M \rightarrow \mathcal{O}_M$ (the Frobenius property).
5. g is flat, i.e., the Levi-Civita connection ∇ associated to g is flat.
6. The $(3, 1)$ -tensor $\nabla \circ$ is totally symmetric (the potentiality condition).
7. $e \in \Gamma(M, \mathcal{T}_M)$ is an identity element for $\circ$.

8. e is flat, i.e., $\nabla e = 0$.

9. $E \in \Gamma(M, \mathcal{T}_M)$ satisfies

$$\nabla \nabla E = 0, \quad (6)$$

$$\text{Lie}_E(\circ) = \circ, \quad (7)$$

and there exists a complex number D such that

$$\text{Lie}_E(g) = Dg. \quad (8)$$

Here, $\mathcal{O}_M$ is the structure sheaf of M , $\mathcal{T}_M$ is the tangent sheaf of M , and Lie_E is the Lie derivative with respect to the vector field E . We call g the metric, E the Euler vector field, and D the charge of the Frobenius manifold M .

Let N denote the dimension of M . Since g is flat, there exists a local coordinate $\{t_a\}_{a=0}^{N-1}$ on M , called the *flat coordinate*, satisfying

$$g(\partial_a, \partial_b) = \eta_{ab}$$

which is unique up to affine transformations. Here, η_{ab} is a constant matrix and

$$\partial_a = \frac{\partial}{\partial t_a}.$$

Since the identity vector field e is flat, we can choose the flat coordinate $\{t_a\}_{a=0}^{N-1}$ so that

$$e = \partial_0.$$

Let Φ_{ab}^c be the structure constant of the product $\circ$ in the flat coordinate:

$$\partial_a \circ \partial_b = \sum_{c=0}^{N-1} \Phi_{ab}^c \partial_c.$$

Then the potentiality condition is that

$$\sum_{e=0}^{N-1} \eta_{ae} \partial_b \Phi_{cd}^e$$

is totally symmetric with respect to a, b, c, d . From this condition, it follows that locally on M , there exists a holomorphic function Φ such that

$$\sum_{d=0}^{N-1} \eta_{ad} \Phi_{bc}^d = \partial_a \partial_b \partial_c \Phi. \quad (9)$$

This function Φ is called the potential of the Frobenius manifold M . Note that

$$\eta_{ab} = \partial_0 \partial_a \partial_b \Phi$$

follows from the Frobenius property.

The associativity of $\circ$ gives the following *WDVV (Witten-Dijkgraaf-Verlinde-Verlinde) equation*

$$\sum_{e,f=0}^{N-1} \frac{\partial^3 \Phi}{\partial t_a \partial t_b \partial t_e} \eta^{ef} \frac{\partial^3 \Phi}{\partial t_f \partial t_c \partial t_d} = \sum_{e,f=0}^{N-1} \frac{\partial^3 \Phi}{\partial t_a \partial t_d \partial t_e} \eta^{ef} \frac{\partial^3 \Phi}{\partial t_f \partial t_b \partial t_c}, \quad (10)$$

which is a non-linear partial differential equation for the potential Φ . Here, η^{ab} is the inverse matrix of $\eta_{ab} : \sum_{c=0}^{N-1} \eta^{ac} \eta_{cb} = \delta_b^a$.

Now let us define $\mathcal{U}, \mathcal{V} \in \Gamma(\mathcal{E}nd(\mathcal{T}_M))$ by

$$\mathcal{U}(X) = E \circ X, \quad (11)$$

$$\mathcal{V}(X) = \nabla_X E - \frac{D}{2} X, \quad (12)$$

and let $\pi : \mathbb{C}^\times \times M \rightarrow M$ be the second projection. Define a connection

$$\widehat{\nabla} : \pi^* \mathcal{T}_M \rightarrow \pi^* \mathcal{T}_M \otimes \Omega_{\mathbb{C}^\times \times M}^1$$

on $\pi^* \mathcal{T}_M$, where $\Omega_{\mathbb{C}^\times \times M}^1$ denotes the cotangent sheaf of $\mathbb{C}^\times \times M$, by

$$\widehat{\nabla}_X Y = \nabla_X Y - \frac{1}{\hbar} X \circ Y, \quad (13)$$

$$\widehat{\nabla}_{\partial_h} Y = \partial_h Y + \frac{1}{\hbar^2} \mathcal{U}(Y) - \frac{1}{\hbar} \mathcal{V}(Y), \quad (14)$$

for X and Y which are pull-backs of local sections of $\mathcal{T}_M$ to $\mathbb{C}^\times \times M$ and $\hbar$ is the inhomogeneous coordinate of $\mathbb{P}^1$, and extend it by the Leibniz rule. Note the $\hbar$ here is the notation by Givental and corresponds to $1/z$ in Dubrovin's papers [9], [11]. $\widehat{\nabla}$ is called the *first structure connection* of the Frobenius manifold M . The following Theorem 5 is a keystone in the theory of Frobenius manifolds:

Theorem 5. $\widehat{\nabla}$ is flat.

For the proof, see the references at the beginning of this section. One can see from the proof that the flatness of $\widehat{\nabla}$ encodes many of the axioms of Frobenius manifolds.

(14) can be considered as a family of meromorphic connections on $\mathbb{P}^1$ parametrized by M , and the flatness of $\widehat{\nabla}$ means that this family is isomonodromic. One can see from (14) that this meromorphic connection has a regular singularity at infinity and an irregular singularity of Poincaré rank 1 at the origin.

Definition 6. A point p on a Frobenius manifold M is semisimple if the product structure $\circ$ on the tangent space $T_p M$ of M at p is semisimple (i.e., if there are no nilpotent elements).

By Dubrovin [9], there exists a local coordinate $\{u_i\}_{i=1}^N$ satisfying

$$\partial_i \circ \partial_j = \delta_{ij} \partial_i, \quad (15)$$

$$E = \sum_{i=1}^N u_i \partial_i, \quad (16)$$

where $\partial_i = \frac{\partial}{\partial u_i}$, in the neighborhood of a semisimple point. Since

$$\begin{aligned} g(\partial_i, \partial_i) &= g(\partial_i, \partial_j \circ \partial_j) \\ &= g(\partial_i \circ \partial_j, \partial_j), \end{aligned}$$

∂_i and ∂_j are orthogonal for $i \neq j$. This coordinate $\{u_i\}_{i=1}^N$ is called the *canonical coordinate*, which is unique up to permutations. By (15) and (16), $\{u_i\}_{i=1}^N$ is characterized as the set of eigenvalues of $\mathcal{U}$.

Let

$$f_i = \Delta_i^{-1/2} \partial_i, \quad i = 1, \dots, N,$$

be the normalized canonical vector field, where

$$\Delta_i = g(\partial_i, \partial_i),$$

and define an $N \times N$ matrix Ψ_{ai} by

$$f_i = \sum_{a=1}^N \Psi_{ai} \frac{\partial}{\partial t_a}.$$

Ψ_{ai} is the coordinate transformation matrix from the normalized canonical coordinate to the flat coordinate. The operator $\mathcal{U}$ defined in (11) is diagonal in the normalized canonical coordinate:

$$\mathcal{U}(\partial_i) = \sum_{j=1}^N u_j \partial_j \circ \partial_i = u_i \partial_i.$$

Let U, V be the matrices representing the operators $\mathcal{U}, \mathcal{V}$ in the normalized canonical coordinate. Then

$$U = \text{diag}(u_1, \dots, u_N)$$

and (14) becomes

$$\partial_{\hbar} Y + \frac{1}{\hbar^2} UY - \frac{1}{\hbar} VY = 0. \quad (17)$$

By Dubrovin [11], Lemma 4.3., there exists a unique $N \times N$ -matrix-valued formal series

$$R(\hbar) = 1 + R_1 \hbar + R_2 \hbar^2 + \dots$$

satisfying

$$R^t(\hbar) R(-\hbar) = 1$$

such that

$$Y = R(\hbar) \tilde{Y}$$

transforms (17) to

$$\hbar \partial_{\hbar} \tilde{Y} + U \tilde{Y} = 0. \quad (18)$$

Here, $\bullet^t$ denotes the transpose of a matrix. Since (18) has a fundamental solution of the form

$$\tilde{Y} = \exp[U/\hbar],$$

(14) has a formal fundamental solution of the form

$$\Phi_{\text{formal}} = \Psi R(\hbar) \exp[U/\hbar] \quad (19)$$

in the flat coordinate.

Definition 7. For $0 \leq \phi < \pi$, a straight line $l = \{\hbar \in \mathbb{C}^\times \mid \arg(\hbar) = \phi, \phi - \pi\}$ passing through the origin is called *admissible* if the line through u_i and u_j is not orthogonal to l for any $i \neq j$.

Fix such a line, and choose a small enough number $\epsilon > 0$ so that any line passing through the origin with angle between $\phi - \epsilon$ and $\phi + \epsilon$ is admissible. Define angular domains D_{right} , D_{left} and D_- by

$$\begin{aligned} D_{\text{right}} &= \{\hbar \in \mathbb{C}^\times \mid \phi - \pi - \epsilon < \arg(\hbar) < \phi + \epsilon\}, \\ D_{\text{left}} &= \{\hbar \in \mathbb{C}^\times \mid \phi - \epsilon < \arg(\hbar) < \phi + \pi + \epsilon\}, \\ D_- &= \{\hbar \in \mathbb{C}^\times \mid \phi - \pi - \epsilon < \arg(\hbar) < \phi - \pi + \epsilon\}. \end{aligned} \quad (20)$$

See Figure 2. Since the singularity at the origin is irregular, the formal solution $\Phi_{\text{formal}}(\hbar)$ does not converge. Nevertheless, by [11] Theorem 4.2., there exist unique solutions $\Phi_{\text{right}}(\hbar)$ and $\Phi_{\text{left}}(\hbar)$, defined on the angular domains D_{right} and D_{left} respectively, which asymptote to the same formal solution:

$$\Phi_{\text{right/left}} \sim \Phi_{\text{formal}} \quad \text{as } \hbar \rightarrow 0 \text{ in } D_{\text{right/left}}.$$




Figure 1: An admissible line and angular domains

Since these two solutions satisfy the same linear differential equation on D_- , there exists a matrix S independent of $\hbar$ such that

$$\Phi_{\text{left}}(\hbar) = \Phi_{\text{right}}(\hbar)S, \quad \hbar \in D_-.$$

This matrix S is called the Stokes matrix.

3 Gromov-Witten invariants

Let X be a smooth projective variety over $\mathbb{C}$. For simplicity, let us assume that $H^k(X; \mathbb{C}) = 0$ for odd k .

Definition 8. $(f, C, p_1, \dots, p_n)$ is a stable map to X of genus g with n marked points if C is a complete curve of arithmetic genus g with at worst ordinary double points, $(p_1, \dots, p_n)$ is an ordered set of n points on C , $f : C \rightarrow X$ is a regular map and the automorphism group of f is finite. In this case, $f_*[C] \in H_2(X, \mathbb{Z})$ is called the degree of f .

Let $\overline{\mathcal{M}}_{g,n}(X; \mathbf{d})$ be the moduli space of stable maps to X of degree $\mathbf{d}$ and

genus g with n marked points. We have the evaluation map

$$\begin{array}{ccc} \text{ev}_i : & \overline{\mathcal{M}}_{g,n}(X; \mathbf{d}) & \longrightarrow X \\ & \cup & \cup \\ & (f, C, p_1, \dots, p_n) & \longmapsto f(p_i). \end{array}$$

Let $\mathcal{L}_i$ be the line bundle on $\overline{\mathcal{M}}_{g,n}(X; \mathbf{d})$ whose fiber over $(f, C, p_1, \dots, p_n) \in \overline{\mathcal{M}}_{g,n}(X; \mathbf{d})$ is the cotangent space $T_{p_i}^*C$ at the i -th marked point p_i . The virtual dimension of $\overline{\mathcal{M}}_{g,n}(X; \mathbf{d})$ is

$$(1 - g)(\dim_{\mathbb{C}} X - 3) + \int_{\mathbf{d}} c_1(TX) + n,$$

where $c_1(TX)$ is the first Chern class of the tangent bundle of X . Given n cohomology classes $\gamma_1, \dots, \gamma_n \in H^*(X; \mathbb{C})$ and non-negative integers $d_1, \dots, d_n$, the Gromov-Witten invariant is defined by

$$\langle \tau_{d_1} \gamma_1, \dots, \tau_{d_n} \gamma_n \rangle_{g, \mathbf{d}} = \int_{[\overline{\mathcal{M}}_{g,n}(X; \mathbf{d})]^{\text{virt}}} \prod_{i=1}^n (c_1(\mathcal{L}_i)^{d_i} \cup \text{ev}_i^*(\gamma_i)).$$

Here, $[\overline{\mathcal{M}}_{g,n}(X; \mathbf{d})]^{\text{virt}}$ is the *virtual fundamental class* [3], [24].

Let $\{T_a\}_{a=0}^{N-1}$ be a basis of $H^*(X; \mathbb{C})$ and let $\{t_a\}_{a=0}^{N-1}$ be the corresponding coordinate of $H^*(X; \mathbb{C})$. Set $\deg t_a = \deg T_a = k$ when $T_a \in H^k(X; \mathbb{C})$. Assume that the basis $\{T_a\}_{a=0}^{N-1}$ are taken so that T_0 is the basis of $H^0(X; \mathbb{C})$. Let

$$\gamma = \sum_{a=0}^{N-1} t_a T_a.$$

The following formal power series $\Phi(\gamma)$ in $\{t_i\}_{i=0}^{N-1}$ is called the *Gromov-Witten potential*:

$$\Phi(\gamma) = \sum_{n=0}^{\infty} \sum_{\mathbf{d} \in H_2(X, \mathbb{Z})} \frac{1}{n!} \langle \underbrace{\gamma, \dots, \gamma}_{n \text{ times}} \rangle_{0, \mathbf{d}}.$$

Since

$$\frac{1}{n!} \langle \underbrace{\gamma, \dots, \gamma}_{n \text{ times}} \rangle_{0, \mathbf{d}} = \sum_{|\alpha|=n} \langle \underbrace{T_0, \dots, T_0}_{\alpha_0 \text{ times}}, \dots, \underbrace{T_{N-1}, \dots, T_{N-1}}_{\alpha_{N-1} \text{ times}} \rangle_{0, \mathbf{d}} \frac{t^\alpha}{\alpha!}$$

where $\alpha = (\alpha_0, \dots, \alpha_{N-1})$ is a multi-index, $|\alpha| = \alpha_0 + \dots + \alpha_{N-1}$, $\alpha! = \alpha_0! \dots \alpha_{N-1}!$ and $t^\alpha = t_0^{\alpha_0} \dots t_{N-1}^{\alpha_{N-1}}$, Φ is the generating function of the Gromov-Witten invariants.

Theorem 9. *When the Gromov-Witten potential Φ converges in some domain D in $H^*(X; \mathbb{C})$, D has a structure of a Frobenius manifold with Φ as a potential function.*

See the references at the beginning of section 2 for details. The above theorem means that although the moduli space of stable maps of genus zero are divided into connected components by the degree and the number of marked points, the Gromov-Witten invariants as a whole have a strong structure so that together they turn $H^*(X; \mathbb{C})$ into a Frobenius manifold. A part of the Frobenius structure is a product structure on the tangent bundle. Since $TH^*(X; \mathbb{C}) \cong H^*(X; \mathbb{C}) \times H^*(X; \mathbb{C})$, it is a product structure on $H^*(X; \mathbb{C})$ parametrized by $H^*(X; \mathbb{C})$ itself. By the *Point Mapping Axiom* in Gromov-Witten theory, this product structure is a deformation of the usual cup product on the cohomology ring, and is called the *quantum cohomology ring*. The metric g is given by the Poincaré pairing on the cohomology group. The unit vector field is

$$e = \frac{\partial}{\partial t_0},$$

the Euler vector field is

$$E = \sum_{a=0}^{N-1} \left(1 - \frac{\deg t_a}{2}\right) t_a \frac{\partial}{\partial t_a} + \sum_{\deg t_b=2} r_b \frac{\partial}{\partial t_b}, \quad (21)$$

and the charge is

$$D = 2 - \dim_{\mathbb{C}} X.$$

Here, $r_b \in \mathbb{Z}$ is defined by

$$c_1(TX) = \sum_{\deg t_b=2} r_b T_b. \quad (22)$$

The WDVV equation (10) is a highly non-trivial differential equation for the Gromov-Witten potential Φ , which allows us to compute all the genus-zero Gromov-Witten invariants from a few initial conditions in special cases. Fro the Gromov-Witten invariants of rational surfaces, see, e.g., [5], [7], [8], [12], [16], [22] and references therein.

4 Mirror symmetry for toric manifolds

In this section, we review the mirror symmetry for toric varieties. See also [2], [13], [15], [20].

Let Σ be a complete fan in $N \cong \mathbb{Z}^n$ and X_Σ be the corresponding toric variety. We denote the primitive generators of one-dimensional cones of Σ by $\{v_k\}_{k=1}^{n+r}$, $v_k = (v_{k1}, \dots, v_{kn}) \in \mathbb{Z}^n$. Here, r is the Picard number of X_Σ . If we put $M = \text{Hom}(N, \mathbb{Z})$, we have an exact sequence

$$0 \rightarrow M \rightarrow \mathbb{Z}^{n+r} \rightarrow H^2(X_\Sigma; \mathbb{Z}) \rightarrow 0,$$

where the homomorphism $M \rightarrow \mathbb{Z}^{n+r}$ is defined by

$$M \ni m \mapsto (m(v_k))_{k=1}^{n+r} \in \mathbb{Z}^{n+r}.$$

Choose a basis $\{T_a\}_{a=1}^r$ of $H^2(X; \mathbb{Z})$ and let $t = (t_1, \dots, t_r)$ be the corresponding coordinate of $H^2(X; \mathbb{Z})$. One-dimensional cones in Σ correspond to line bundles on X_Σ , and let w_k be the first Chern class of the line bundle corresponding to the one-dimensional cone generated by v_k . Define an $(n+r) \times r$ -matrix m_{ka} by

$$w_k = \sum_{a=1}^r m_{ka} T_a. \quad (23)$$

Now, following Givental, introduce an $H^*(X_\Sigma; \mathbb{C})$ -valued formal function $I(t_0, t; \hbar)$ of $t_0 \in H^0(X_\Sigma; \mathbb{C})$, $t \in H^2(X_\Sigma; \mathbb{C})$ and $\hbar \in \mathbb{C}^\times$ by

$$I(t_0, t; \hbar) = e^{(t_0 + Tt)/\hbar} \sum_{\mathbf{d} \in H_2(X_\Sigma; \mathbb{Z})_{\text{eff}}} q^{\mathbf{d}} \prod_{k=1}^{n+r} \frac{\prod_{l=-\infty}^0 (w_k + l\hbar)}{\prod_{l=-\infty}^{\langle \mathbf{d}, w_k \rangle} (w_k + l\hbar)}. \quad (24)$$

Here, $Tt = T_1 t_1 + \dots + T_r t_r$, $H_2(X_\Sigma; \mathbb{Z})_{\text{eff}} \subset H_2(X_\Sigma; \mathbb{Z})$ is the cone spanned by the homology class of curves and

$$q^{\mathbf{d}} = q_1^{\langle \mathbf{d}, T_1 \rangle} \dots q_r^{\langle \mathbf{d}, T_r \rangle},$$

where $q_a = \exp(t_a)$, $a = 1, \dots, r$ and $\langle \bullet, \bullet \rangle$ is the pairing between homology and cohomology.

It follows from (24) that the I -function satisfies the following set of hypergeometric equations of the GKZ (Gelfand-Kapranov-Zelevinski) type:

$$D_{\mathbf{d}} I = 0, \quad \mathbf{d} \in H_2(X_\Sigma; \mathbb{Z}), \quad (25)$$

where

$$D_{\mathbf{d}} = q^{\mathbf{d}} \prod_{\langle w_k, \mathbf{d} \rangle < 0} \prod_{l=0}^{-\langle w_k, \mathbf{d} \rangle - 1} U_{kl} - \prod_{\langle w_k, \mathbf{d} \rangle > 0} \prod_{l=0}^{\langle w_k, \mathbf{d} \rangle - 1} U_{kl},$$

$$U_{kl} = \sum_{a=1}^r m_{ka} \hbar \frac{\partial}{\partial t_a} - l\hbar.$$

From now on, we assume that the anticanonical bundle $-K_{X_\Sigma}$ of the toric variety X_Σ is nef. Note that although I -function is an infinite series in both positive and negative powers of $\hbar$ for general X_Σ , it contains only non-positive powers of $\hbar$ when $-K_{X_\Sigma}$ is nef. This follows from the fact that $\exp[-(t_0 + Tt)/\hbar]I$ is homogeneous of degree zero with respect to the degree assigned as

$$\deg w_i = \deg \hbar = 1, \quad \deg(q^{\mathbf{d}}) = \langle \mathbf{d}, c_1(TX_\Sigma) \rangle. \quad (26)$$

Therefore, the coefficient of a positive power of $\hbar$ must have a negative degree, which is impossible when $-K_{X_\Sigma}$ is nef.

I learned the following Lemma 10 from H. Iritani:

Lemma 10. *When the anticanonical bundle $-K_{X_\Sigma}$ of a toric variety X_Σ is nef, there exists a domain $D \subset H^2(X_\Sigma; \mathbb{C})$ such that $I(t_0, t; \hbar)$ converges for any $t_0 \in H^0(X_\Sigma; \mathbb{C})$, $t \in D$ and $|\hbar| > 1$*

Proof. Fix a norm $\|\bullet\|$ on $H^*(X_\Sigma; \mathbb{C})$ such that $\|\omega\tau\| \leq \|\omega\|\|\tau\|$. Let C_1 be a positive number satisfying

$$\sup_{|\hbar|=1, l \geq 1, k=1, \dots, n+r} \left(\|w_k\|, \left\| 1 + \frac{w_k}{l\hbar} \right\|, \left\| \left(1 + \frac{w_k}{l\hbar} \right)^{-1} \right\| \right) \leq C_1.$$

Then for $\hbar \in \mathbb{C}$ such that $|\hbar| = 1$,

$$\begin{aligned} \left\| \prod_{k=1}^{n+r} \frac{\prod_{l=-\infty}^0 (w_k + l\hbar)}{\prod_{l=-\infty}^{\langle \mathbf{d}, w_k \rangle} (w_k + l\hbar)} \right\| &\leq \frac{\prod_{\langle \mathbf{d}, w_k \rangle < 0} |\langle \mathbf{d}, w_k \rangle|!}{\prod_{\langle \mathbf{d}, w_k \rangle > 0} \langle \mathbf{d}, w_k \rangle!} C_1^{\sum_{k=1}^{n+r} |\langle \mathbf{d}, w_k \rangle|} \\ &\leq \frac{C_2^{|\mathbf{d}|}}{\langle \mathbf{d}, c_1(X) \rangle!} \end{aligned}$$

Here, C_2 is some positive number and $|\mathbf{d}| = |\langle \mathbf{d}, T_1 \rangle| + \dots + |\langle \mathbf{d}, T_r \rangle|$. The second inequality follows from

$$\begin{aligned} n_1! n_2! \cdots n_k! &\leq (n_1 + \cdots + n_k)!, \\ \frac{1}{n_1! n_2! \cdots n_k!} &= \frac{1}{(n_1 + \cdots + n_k)!} \binom{n_1 + \cdots + n_k}{n_1, \dots, n_k} \\ &\leq \frac{1}{(n_1 + \cdots + n_k)!} k^{n_1 + \cdots + n_k}, \\ \frac{n!}{m!} &\leq \frac{1}{(m-n)!} \quad \text{if } n \leq m. \end{aligned}$$

and

$$\sum_{k=1}^{n+r} \langle \mathbf{d}, w_k \rangle = \langle \mathbf{d}, -K_{X_\Sigma} \rangle \geq 0$$

since $\mathbf{d}$ is an effective class and $-K_{X_\Sigma}$ is nef. Therefore, if we define $D = \{t \in H^2(X_\Sigma; \mathbb{C}) \mid q_a < 1/C_2, a = 1, \dots, r\}$, the I -function converges when $|\hbar| = 1$ and $t \in D$. Since the I -function contains only non-positive powers of $\hbar$, it also converges when $|\hbar| > 1$ and $t \in D$. $\square$

Now, let us introduce another $H^*(X_\Sigma; \mathbb{C})$ -valued formal function $J(t_0, t; \hbar)$ of $t_0 \in H^0(X_\Sigma; \mathbb{C})$, $t \in H^2(X_\Sigma; \mathbb{C})$ and $\hbar \in \mathbb{C}^\times$ as follows: First, define a set of $H^*(X_\Sigma; \mathbb{C})$ -valued formal functions $\{s_a\}_{a=0}^{N-1}$ of $\{t_a\}_{a=0}^r$ by

$$s_a = T_a + \sum_{n=0}^{\infty} \hbar^{-n-1} \sum_{b=0}^{N-1} \sum_{k=0}^{\infty} \sum_{\mathbf{d} \in H_2(X_\Sigma; \mathbb{C})} \frac{1}{k!} \langle \tau_n T_a, T_b, \underbrace{\gamma, \dots, \gamma}_{k \text{ times}} \rangle_{0, \mathbf{d}} T^b,$$

where

$$\gamma = \sum_{a=0}^r t_a T_a.$$

Here, T^a is the Poincaré dual of T_a :

$$g(T_a, T^b) = \delta_{ab}.$$

Then it follows from the *topological recursion relation* that

$$\hbar \frac{\partial s_a}{\partial t_b} = T_b \circ s_a, \quad b = 0, 1, \dots, r. \quad (27)$$

See, e.g., [6] Proposition 10.2.1. Note that (27) is the t_b -part of the differential equation for flat sections of the first structure connection (13). Now define the *Givental's J-function* by

$$J = \sum_{a=0}^{N-1} g(s_a, 1) T^a.$$

Since $H^*(X_\Sigma; \mathbb{C})$ is generated by $H^2(X_\Sigma; \mathbb{C})$ as a ring, there exists a set of polynomials $\{P_a(x_1, \dots, x_r)\}_{a=0}^{N-1}$ of r variables such that

$$T_a = P_a(T_1, \dots, T_r), \quad a = 0, 1, \dots, N-1.$$

Since

$$T_a \circ T_b = T_a \cup T_b + O(q)$$

by the Point Mapping Axiom, there exists a set $\{\tilde{P}_a(x_1, \dots, x_r)\}_{a=0}^{N-1}$ of polynomials whose coefficient is a formal power series in q such that

$$\tilde{P}_a(\hbar \frac{\partial}{\partial t_1} + T_1 \circ, \dots, \hbar \frac{\partial}{\partial t_r} + T_r \circ) 1 = T_a, \quad a = 0, 1, \dots, N-1.$$

Then, since

$$P(\hbar \partial_1, \dots, \hbar \partial_r) J = \sum_{a=0}^{N-1} g \left(s_a, P(\hbar \frac{\partial}{\partial t_1} + T_1 \circ, \dots, \hbar \frac{\partial}{\partial t_r} + T_r \circ) 1 \right) T^a$$

for a polynomial $P(x_1, \dots, x_r)$,

$$s_a = \sum_{b=0}^{N-1} g \left(\tilde{P}_b(\hbar \partial_1, \dots, \hbar \partial_r) J, T_a \right) T^b.$$

To sum up, we have introduced two $H^*(X_\Sigma; \mathbb{C})$ -valued formal functions $I(t_0, t; \hbar)$ and $J(t_0, t; \hbar)$. The I -function is defined in terms of the combinatorial data of the fan Σ defining X_Σ and satisfies a set of hypergeometric differential equations (25) of GKZ type. The J -function is defined in terms of the Gromov-Witten invariants of X_Σ and encodes the information of the solution to the equation (27), hence the structure of the product $\circ$.

The mirror theorem by Givental (see [15] Theorem 0.2, Proposition 6.4, and the discussion at the end of section 7) gives the relation between the I -function and the J -function as follows: Since $-K_{X_\Sigma}$ is nef, all the variables q_a 's have non-negative degrees, see (26). It follows from the homogeneity of $\exp[-(t_0 + Tt)/\hbar] I(t_0, t; \hbar)$ that the I -function has the form

$$I(t_0, t; \hbar) = e^{(t_0 + Tt)/\hbar} \left(I^{(0)}(q) + (I^{(10)}(q) + I^{(11)}(q)) \hbar^{-1} + O(\hbar^{-2}) \right), \quad (28)$$

where $I^{(0)}(q)$ and $I^{(10)}(q)$ are $H^0(X_\Sigma; \mathbb{C})$ -valued and $I^{(11)}(q)$ is $H^2(X_\Sigma; \mathbb{C})$ -valued. Then, the J -function is given by

$$\begin{aligned} J(\tilde{t}_0, \tilde{t}; \hbar) &= I(t_0, t; \hbar) / I^{(0)}(q) \\ &= \exp[(t_0 + Tt)/\hbar] \left(1 + I^{(0)}(q)^{-1} (I^{(10)}(q) + I^{(11)}(q)) \hbar^{-1} + O(\hbar^{-2}) \right) \\ &= \exp \left[\left((t_0 + I^{(0)}(q)^{-1} I^{(10)}(q)) + (t + I^{(0)}(q)^{-1} I^{(11)}(q)) \right) / \hbar \right] (1 + O(\hbar^{-2})) \\ &= \exp \left[(\tilde{t}_0 + T\tilde{t}) / \hbar \right] (1 + O(\hbar^{-2})). \end{aligned}$$

Here, we have performed a homogeneous coordinate transformation

$$\tilde{t}_0 = t_0 + I^{(0)}(q)^{-1} I^{(10)}(q), \quad \tilde{t} = t + I^{(0)}(q)^{-1} I^{(11)}(q),$$

called the *mirror transformation*. Since the I -function is convergent, the mirror transformation and the J -function are also convergent.

5 Stationary-phase integral and Stokes matrix

Let $v_0 = (0, \dots, 0) \in \mathbb{Z}^n$, and $\{v_k\}_{k=1}^{n+r} \subset \mathbb{Z}^n$ be the set of primitive generators of one-dimensional cones of the fan Σ in $\mathbb{Z}^n$ defining a toric variety X_Σ as in section 4. We assume that the anticanonical bundle $-K_{X_\Sigma}$ of X_Σ is nef. Define a Laurent polynomial W by

$$W(x_1, \dots, x_n) = \sum_{k=0}^{n+r} c_k x_1^{v_{k1}} \cdots x_r^{v_{kr}}. \quad (29)$$

W defines a regular function on the algebraic torus $(\mathbb{C}^\times)^n$. By Kouchnirenko [23], the number N of critical points of W is $n!$ times the volume of the Newton polygon of W (i.e., the convex hull of $\{v_k\}_{k=1}^{n+r}$ when $c_k \in \mathbb{C}$, $k = 0, \dots, n+r$, are general enough. This N is equal to the rank of the total cohomology ring $H^*(X_\Sigma, \mathbb{C})$ of X_Σ . Let $\{p_i\}_{i=1}^N$ be the set of critical points of W . They are functions of $\{c_k\}_{k=0}^{n+r}$. Fix a complete Kähler metric on the algebraic torus $(\mathbb{C}^\times)^n$. For a general $\hbar$, let $\Gamma_i(\hbar)$ be the descending Morse cycle for $\Re(W/\hbar)$. The image of $\Gamma_i(\hbar)$ by W is a half-line starting from the critical value $W(p_i)$, and the fiber above a point p on this half-line is the cycle in $W^{-1}(p)$ which vanishes at p_i by the parallel transport along this half-line. Following Givental [14], for $i = 1, \dots, N$, consider the integral

$$\mathcal{I}_i = (\pi\hbar)^{-n/2} \int_{\Gamma_i(\hbar)} e^{\frac{1}{\hbar}W(x)} \frac{dx_1 \wedge \cdots \wedge dx_n}{x_1 \cdots x_n}. \quad (30)$$

This integral is invariant under the natural action of the torus $(\mathbb{C}^\times)^n$ on itself defined by

$$(\mathbb{C}^\times)^n \ni (\alpha_1, \dots, \alpha_n) : (x_1, \dots, x_n) \mapsto (\alpha_1 x_1, \dots, \alpha_n x_n).$$

Therefore, if we define the variables $\{t_a\}_{a=0}^r$ by $t_0 = c_0$ and

$$\exp(t_a) = c_1^{m_{1a}} \cdots c_{n+r}^{m_{n+r,a}}, \quad a = 1, \dots, r,$$

where m_{ka} 's are defined in (23), $\mathcal{I}_i$'s depend on $\{c_k\}_{k=0}^{n+r}$ only through $\{t_a\}_{a=0}^r$.

It is easy to see that $\mathcal{I}_i$'s satisfy the same differential equations (25) as the I -function. Since the solution space to these differential equation is an N -dimensional $\mathbb{C}$ -vector space and $\{\mathcal{I}_i\}_{i=1}^N$ are linearly independent, the set of components of the I -function and $\{\mathcal{I}_i\}_{i=1}^N$ differ only by an $\hbar$ -dependent linear transformation.

The $\hbar$ -dependence of $\mathcal{I}_i$ is simple. If we assign degree 1 to variables $t_0, \hbar$ and $\{x_i\}_{i=1}^n$, and degree r_a to the variable $\exp(t_a)$ for $a = 1, \dots, r$ where r_a 's are defined in (22), $\mathcal{I}_i$ is homogeneous of degree $-n/2$:

$$\left[\hbar \frac{\partial}{\partial \hbar} + t_0 \frac{\partial}{\partial t_0} + \sum_{a=1}^r r_a \frac{\partial}{\partial t_a} + \frac{n}{2} \right] \mathcal{I}_i = 0. \quad (31)$$

The important point is that the $\hbar$ -part (14) of the differential equation for the $H^{2n}(X_\Sigma; \mathbb{C})$ -component of a flat section of the first structure connection is precisely the homogeneity condition (31): Assume that a section Y of $\pi^* \mathcal{T}_M$ satisfies

$$\widehat{\nabla}_X Y = \nabla_X Y - \frac{1}{\hbar} X \circ Y = 0$$

for any section X of $\pi^* T_M$. Then

$$\begin{aligned} \hbar \widehat{\nabla}_{\partial_\hbar} Y &= \hbar \partial_\hbar Y + \frac{1}{\hbar} \mathcal{U}(Y) - \mathcal{V}(Y) \\ &= \hbar \partial_\hbar Y + \frac{1}{\hbar} E \circ Y - \mathcal{V}(Y) \\ &= \hbar \partial_\hbar Y + \nabla_E Y - \mathcal{V}(Y). \end{aligned}$$

Note that for the flat coordinates $\{t_a\}_{a=0}^{N-1}$,

$$\begin{aligned} \mathcal{V}(\partial_a) &= \nabla_{\partial_a} E - \frac{D}{2} \partial_a \\ &= \nabla_{\partial_a} \left(\sum_{b=0}^{N-1} \left(1 - \frac{\deg t_b}{2} \right) t_b \partial_b + \sum_{\deg t_c=2} r_c \partial_c \right) - \frac{2-n}{2} \partial_a \\ &= \left(\frac{n - \deg t_a}{2} \right) \partial_a. \end{aligned}$$

Since ∇_E counts the degree, this differential equation requires the $H^k(X_\Sigma; \mathbb{C})$ -component of the flat section of the first structure connection to be homogeneous of degree $(n - k)/2$.

Now, recall that by Givental's mirror theorem, $I^{(0)}(q)^{-1} I(t_0, t; \hbar)$ coincides with the J -function, which have the same dependence on $\{t_a\}_{a=0}^r$ with the $H^{2n}(X_\Sigma; \mathbb{C})$ -components of the flat section for the first structure connection. Then, since the stationary-phase integrals (30) have the same dependence on $\{t_a\}_{a=0}^r$ as the I -function and satisfies the desired homogeneity condition (31), $I^{(0)}(q)^{-1} \mathcal{I}_i(t_0, t; \hbar)$ gives the $H^{2n}(X_\Sigma; \mathbb{C})$ -component of a flat section for the first structure connection for $i = 1, \dots, N$.

By the saddle-point approximation,

$$I^{(0)}(q)^{-1}\mathcal{I}_i \sim \frac{I^{(0)}(q)^{-1}}{p_{i,1} \cdots p_{i,n}} \left[\det \left(\frac{\partial^2 W}{\partial x_l \partial x_m}(p_i) \right)_{l,m} \right]^{-\frac{1}{2}} (1 + O(\hbar)) e^{\frac{1}{\hbar}W(p_i)} \quad (32)$$

as $|\hbar| \rightarrow 0$ with $\arg \hbar$ fixed, where $p_i = (p_{i,1}, \dots, p_{i,n}) \in (\mathbb{C}^\times)^n$ is the i -th critical point of W . This gives the asymptotic expansion appearing in (19).

Since the Stokes matrix acts on all the components of the fundamental solution equally, it is enough to compute its action on the $H^{2n}(X_\Sigma; \mathbb{C})$ -component computed above. The integration cycle $\Gamma_i(\hbar)$ in (30) undergoes a discontinuous change when $\hbar$ crosses the line such that the half-line starting from the critical value u_i in the direction of $-\hbar^{-1}$ passes through another critical value u_j . Therefore, $\mathcal{I}_i$ is not holomorphic on such a line. Since $\mathcal{I}_i$ has a monodromy, it is impossible to obtain a holomorphic function on the whole $\hbar$ -plane by analytic continuation, but one can obtain a holomorphic function on the two angular domains D_{left} and D_{right} defined in Section 2 as follows: Consider the local system on $\mathbb{C}^\times$ whose fiber over $\hbar \in \mathbb{C}^\times$ is the relative homology group $H_n((\mathbb{C}^\times)^n, \Re(W/\hbar) \ll 0; \mathbb{Z})$. Define two sets of sections $\{\Gamma_{i,\text{left}}(\hbar)\}_{i=1}^N$ and $\{\Gamma_{i,\text{right}}(\hbar)\}_{i=1}^N$ of this local system on D_{left} and D_{right} by the condition that they coincide with the classes defined by the descending Morse cycles at $\hbar = \exp[(\phi + \pi/2)\sqrt{-1}]$ and $\hbar = \exp[(\phi - \pi/2)\sqrt{-1}]$ respectively;

$$\begin{aligned} \Gamma_{i,\text{left}}(\exp[(\phi + \pi/2)\sqrt{-1}]) &= \Gamma_i(\exp[(\phi + \pi/2)\sqrt{-1}]), \\ \Gamma_{i,\text{right}}(\exp[(\phi - \pi/2)\sqrt{-1}]) &= \Gamma_i(\exp[(\phi - \pi/2)\sqrt{-1}]). \end{aligned}$$

Since D_{left} and D_{right} are simply-connected, these conditions determine $\Gamma_{i,\text{left}}(\hbar)$ and $\Gamma_{i,\text{right}}(\hbar)$ uniquely. Define $\mathcal{I}_{i,\text{left}}$ (resp. $\mathcal{I}_{i,\text{right}}$) by the integral (30) with $\Gamma_{i,\text{left}}(\hbar)$ (resp. $\Gamma_{i,\text{right}}(\hbar)$) as the integration cycle. Then $\mathcal{I}_{i,\text{left}}$ (resp. $\mathcal{I}_{i,\text{right}}$) are holomorphic and have the asymptotic behaviors (32) as $\hbar \rightarrow 0$ in D_{left} (resp. D_{right}).

Figure 2 shows the images Γ_i 's of $\Gamma_{i,\text{right}}(\hbar)$'s by W at $\hbar = 1$. Here, we have assumed that the imaginary axis $\Re(\hbar) = 0$ is admissible, and have chosen it as an admissible line l . Figure 3 shows the images $\Gamma_{i,l}$'s and $\Gamma_{i,r}$'s of $\Gamma_{i,\text{left}}(\hbar)$'s and $\Gamma_{i,\text{right}}(\hbar)$'s by W at $\hbar = -\sqrt{-1}$.

Since the integrand is single-valued, the Stokes matrix is given by the transformation matrix between these two sets of cycles. To describe it, let us first choose a regular value $p \in \mathbb{C}$ such that $\Re(p)$ is small enough so that the line segment c_i from the i -th critical value u_i to p is above c_j for any $j > i$ (i.e., if $x \in c_i$, $y \in c_j$ and $\Re(x) = \Re(y)$, then $\Im(x) > \Im(y)$). Then, the ordered set of cycles $(C_i)_{i=1}^N$ in $W^{-1}(p)$, which vanish along the paths $(c_i)_{i=1}^N$




Figure 2: Integration cycles at $\hbar = 1$ Figure 3: Integration cycles at $\hbar = -\sqrt{-1}$

respectively, forms a *distinguished basis of vanishing cycles*, see e.g. [1]. It is obvious that

$$\Gamma_{1,\text{left}}(\hbar) = \Gamma_{1,\text{right}}(\hbar).$$

in $H_n((\mathbb{C}^\times)^n, \Re[W/\hbar] \ll 0; \mathbb{Z})$ in the neighborhood of $\hbar = -\sqrt{-1}$. As for $\Gamma_2(\hbar)$, it follows from the Picard-Lefschetz formula that

$$\Gamma_{2,\text{left}}(\hbar) = \Gamma_{2,\text{right}}(\hbar) - (C_1, C_2)\Gamma_{1,\text{right}}(\hbar),$$

where $(\bullet, \bullet)$ is the intersection form in $H_n(W^{-1}(p); \mathbb{Z})$. In the same way, we have

$$\Gamma_{i,\text{left}}(\hbar) = \Gamma_{i,\text{right}}(\hbar) - (C_{i-1}, C_i)\Gamma_{i-1,\text{right}}(\hbar) - \cdots - (C_1, C_i)\Gamma_{1,\text{right}}(\hbar),$$

by successive use of the Picard-Lefschetz formula, from which

$$S_{ij} = \begin{cases} 1 & \text{if } i = j, \\ -(C_i, C_j) & \text{if } i < j, \\ 0 & \text{otherwise} \end{cases} \quad (33)$$

follows.

6 Intersection numbers of vanishing cycles

Now we consider the toric variety Y which is the projective plane blown-up at 6 points including infinitely-near points. Figure 4 shows the generators of

the one-dimensional cones of the fan defining Y .




Figure 4: the toric data of Y

Y is deformation-equivalent to cubic surfaces, and $-K_Y$ is not ample but nef. The mirror for Y is defined by (29), which we specialize to $c_3 = c_8 = c_9 = 1$ and $c_i = 0$ for $i \neq 3, 8, 9$, to obtain

$$W(x, y) = \frac{x^2}{y} + \frac{y^2}{x} + \frac{1}{xy}, \quad (34)$$

which is generic in the sense that all the critical points are non-degenerate. In this case, $W^{-1}(0)$ is a Fermat curve in $(\mathbb{C}^\times)^2$:

$$W^{-1}(0) = \{(x, y) \in (\mathbb{C}^\times)^2 \mid x^3 + y^3 + 1 = 0\}.$$

W has nine critical points $\{p_{ij}\}_{i,j=0}^2$, $p_{ij} = (\omega^{i+j-1}, \omega^{2j})$, where we have fixed a primitive cubic root ω of unity. Let C_{ij} be the vanishing cycle in $W^{-1}(0)$ which vanishes at the critical point p_{ij} along the straight line from 0 to the critical value $3\omega^{1-i}$. Then $(C_{00}, C_{01}, C_{02}, C_{10}, C_{11}, C_{12}, C_{20}, C_{21}, C_{22})$ is a distinguished basis of vanishing cycles.

Figure 5 shows these vanishing cycles with orientations chosen by hand. Opposite sides of the square is identified to form a torus, and the circles denote nine points which are missing in $W^{-1}(0)$ since it is not compact. Note that the quotient of $W^{-1}(0)$ by the action of $\mathbb{Z}/(3\mathbb{Z})$ generated by

$$(x, y) \mapsto (\omega x, \omega^2 y).$$

is

$$\left\{ (X, Y) \in (\mathbb{C}^\times)^2 \mid X + Y + \frac{1}{XY} = 0 \right\}$$

appearing in Seidel's work [26]. This action is given by the translation in the direction of the diagonal line from the lower-left corner to the upper-right




Figure 5: Vanishing cycles in $W^{-1}(0)$

corner of the square in Figure 5 by one-third of the length of the diagonal line, and the quotient by this action is exactly Figure 2 in [26].

Intersection numbers of C_{ij} 's are

$$(C_{ij}, C_{ik}) = 0 \quad \text{if } j \neq k,$$

and

$$(C_{ij}, C_{kl}) = -1 \quad \text{if } i < k.$$

7 Derived category of coherent sheaves

Let $\mathbb{Z}/3\mathbb{Z}$ act on a three-dimensional vector space $V \cong \mathbb{C}^3$ by

$$\mathbb{Z}/3\mathbb{Z} \ni [1] : V \ni (x : y : z) \mapsto (\omega x : \omega^2 y : z) \in V$$

where ω is a primitive cubic root of unity. This action defines an action of $\mathbb{Z}/3\mathbb{Z}$ on $\mathbb{P}^2 = \mathbb{P}(V)$, and the toric variety Y defined in Section 6 is the minimal resolution of the quotient $\mathbb{P}(V)/(\mathbb{Z}/3\mathbb{Z})$. By Beilinson [4], $(\mathcal{E}_0, \mathcal{E}_1, \mathcal{E}_2) = (\mathcal{O}_{\mathbb{P}^2}(-1), \Omega_{\mathbb{P}^2}(1), \mathcal{O}_{\mathbb{P}^2})$ is an exceptional collection generating $D^b \text{coh} \mathbb{P}^2$, and by Kapranov-Vasserot [21], we have

$$D^b \text{coh} Y \cong D^b \text{coh}^{\mathbb{Z}/3\mathbb{Z}} \mathbb{P}^2.$$

Here, $\mathcal{O}_{\mathbb{P}^2}(-1)$ is the tautological sheaf, $\Omega_{\mathbb{P}^2}(1)$ is the cotangent sheaf tensored with the dual of $\mathcal{O}_{\mathbb{P}^2}(-1)$, and $\text{coh}^{\mathbb{Z}/3\mathbb{Z}}\mathbb{P}^2$ is the category of $\mathbb{Z}/3\mathbb{Z}$ -equivariant coherent sheaves on $\mathbb{P}^2$. Now, let $\mathcal{E}_{ij}$ be the object of $D^b\text{coh}Y$ corresponding to the $\mathbb{Z}/3\mathbb{Z}$ -equivariant coherent sheaf $\mathcal{E}_i \otimes \rho_j$ on $\mathbb{P}^2$ by the above equivalence. Here, $\rho_j : \mathbb{Z}/3\mathbb{Z} \ni [1] \mapsto \omega^j \in \mathbb{C}^\times$ is a one-dimensional representation of $\mathbb{Z}/3\mathbb{Z}$. Then $(\mathcal{E}_{00}, \mathcal{E}_{01}, \mathcal{E}_{02}, \mathcal{E}_{10}, \mathcal{E}_{11}, \mathcal{E}_{12}, \mathcal{E}_{20}, \mathcal{E}_{21}, \mathcal{E}_{22})$ is an exceptional collection generating $D^b\text{coh}Y$. The Ext-groups between them can be calculated as follows: First, we can use the exact sequence

$$0 \rightarrow \Omega_{\mathbb{P}^2}(1) \rightarrow V^\vee \otimes \mathcal{O}_{\mathbb{P}^2} \rightarrow \mathcal{O}_{\mathbb{P}^2}(1) \rightarrow 0$$

to compute the Ext-groups between $\mathcal{E}_i$'s:

$$\begin{aligned} \mathbb{R}\text{Hom}(\mathcal{E}_1, \mathcal{E}_2) &= \mathbb{R}\text{Hom}(\mathcal{O}_{\mathbb{P}^2}(-1), \{V^\vee \otimes \mathcal{O}_{\mathbb{P}^2} \rightarrow \mathcal{O}_{\mathbb{P}^2}(1)\}) \\ &= \mathbb{R}\Gamma(\mathcal{O}_{\mathbb{P}^2}(1) \otimes \{V^\vee \otimes \mathcal{O}_{\mathbb{P}^2} \rightarrow \mathcal{O}_{\mathbb{P}^2}(1)\}) \\ &= \mathbb{R}\Gamma(\{V^\vee \otimes \mathcal{O}_{\mathbb{P}^2}(1) \rightarrow \mathcal{O}_{\mathbb{P}^2}(2)\}) \\ &= \{V^\vee \otimes V^\vee \rightarrow \text{Sym}^2 V^\vee\} \\ &= \wedge^2 V^\vee, \\ \mathbb{R}\text{Hom}(\mathcal{E}_1, \mathcal{E}_3) &= \mathbb{R}\text{Hom}(\mathcal{O}_{\mathbb{P}^2}(-1), \mathcal{O}_{\mathbb{P}^2}) \\ &= \mathbb{R}\Gamma(\mathcal{O}_{\mathbb{P}^2}(1)) \\ &= V^\vee, \\ \mathbb{R}\text{Hom}(\mathcal{E}_2, \mathcal{E}_3) &= \mathbb{R}\text{Hom}(\{V^\vee \otimes \mathcal{O}_{\mathbb{P}^2} \rightarrow \mathcal{O}_{\mathbb{P}^2}(1)\}, \mathcal{O}_{\mathbb{P}^2}) \\ &= \mathbb{R}\Gamma(\{\mathcal{O}_{\mathbb{P}^2}(-1) \rightarrow V \otimes \mathcal{O}_{\mathbb{P}^2}\}) \\ &= V. \end{aligned}$$

Here, $\mathbb{R}\text{Hom}$ and $\mathbb{R}\Gamma$ denote the right derived functor of Hom and Γ (taking global sections) respectively. Then the Ext-groups between $\mathcal{E}_{ij}$'s are given by

$$\begin{aligned} \text{Ext}^k(\mathcal{E}_{ij}, \mathcal{E}_{lm}) &= \text{Ext}^k(\mathcal{E}_i \otimes \rho_j, \mathcal{E}_l \otimes \rho_m) \\ &= (\text{Ext}^k(\mathcal{E}_i, \mathcal{E}_l) \otimes \rho_j^\vee \otimes \rho_m)^{\mathbb{Z}/3\mathbb{Z}}, \end{aligned}$$

where $\bullet^{\mathbb{Z}/3\mathbb{Z}}$ denotes the $\mathbb{Z}/3\mathbb{Z}$ -invariant part. Therefore, the dimension of $\text{Ext}^k(\mathcal{E}_{ij}, \mathcal{E}_{lm})$ is non-zero if and only if

$$i = l, j = m \text{ and } k = 0,$$

or

$$j < l \text{ and } k = 0,$$

and the dimensions in all these cases are one. Therefore, we have

$$-(C_{ij}, C_{lm}) = \sum_{k=0}^2 (-1)^k \dim \text{Ext}^k(\mathcal{E}_{ij}, \mathcal{E}_{lm}) \quad \text{if } i \leq l \text{ and } (i, j) \neq (l, m),$$

which, combined with (33), proves Conjecture 1 for Y .

References

- [1] V. I. Arnol'd, S. M. Guseĭn-Zade, and A. N. Varchenko. *Singularities of differentiable maps. Vol. II*, volume 83 of *Monographs in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1988. Monodromy and asymptotics of integrals, Translated from the Russian by Hugh Porteous, Translation revised by the authors and James Montaldi.
- [2] Victor V. Batyrev. Quantum cohomology rings of toric manifolds. *Astérisque*, (218):9–34, 1993. Journées de Géométrie Algébrique d'Orsay (Orsay, 1992).
- [3] K. Behrend and B. Fantechi. The intrinsic normal cone. *Invent. Math.*, 128(1):45–88, 1997.
- [4] A. A. Beĭlinson. Coherent sheaves on $\mathbf{P}^n$ and problems in linear algebra. *Funktsional. Anal. i Prilozhen.*, 12(3):68–69, 1978.
- [5] Lucia Caporaso and Joe Harris. Parameter spaces for curves on surfaces and enumeration of rational curves. *Compositio Math.*, 113(2):155–208, 1998.
- [6] David A. Cox and Sheldon Katz. *Mirror symmetry and algebraic geometry*, volume 68 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999.
- [7] Bruce Crauder and Rick Miranda. Quantum cohomology of rational surfaces. In *The moduli space of curves (Texel Island, 1994)*, volume 129 of *Progr. Math.*, pages 33–80. Birkhäuser Boston, Boston, MA, 1995.
- [8] P. Di Francesco and C. Itzykson. Quantum intersection rings. In *The moduli space of curves (Texel Island, 1994)*, volume 129 of *Progr. Math.*, pages 81–148. Birkhäuser Boston, Boston, MA, 1995.
- [9] Boris Dubrovin. Geometry of 2D topological field theories. In *Integrable systems and quantum groups (Montecatini Terme, 1993)*, volume 1620 of *Lecture Notes in Math.*, pages 120–348. Springer, Berlin, 1996.
- [10] Boris Dubrovin. Geometry and analytic theory of Frobenius manifolds. In *Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998)*, number Extra Vol. II, pages 315–326 (electronic), 1998.
- [11] Boris Dubrovin. Painlevé transcendents in two-dimensional topological field theory. In *The Painlevé property*, CRM Ser. Math. Phys., pages 287–412. Springer, New York, 1999.

- [12] Andreas Gathmann. Gromov-Witten invariants of blow-ups. *J. Algebraic Geom.*, 10(3):399–432, 2001.
- [13] Alexander Givental. Equivariant Gromov-Witten invariants. *Internat. Math. Res. Notices*, (13):613–663, 1996.
- [14] Alexander Givental. Elliptic Gromov-Witten invariants and the generalized mirror conjecture. In *Integrable systems and algebraic geometry (Kobe/Kyoto, 1997)*, pages 107–155. World Sci. Publishing, River Edge, NJ, 1998.
- [15] Alexander Givental. A mirror theorem for toric complete intersections. In *Topological field theory, primitive forms and related topics (Kyoto, 1996)*, volume 160 of *Progr. Math.*, pages 141–175. Birkhäuser Boston, Boston, MA, 1998.
- [16] L. Göttsche and R. Pandharipande. The quantum cohomology of blow-ups of $\mathbf{P}^2$ and enumerative geometry. *J. Differential Geom.*, 48(1):61–90, 1998.
- [17] Davide Guzzetti. Stokes matrices and monodromy of the quantum cohomology of projective spaces. *Comm. Math. Phys.*, 207(2):341–383, 1999.
- [18] Claus Hertling. *Frobenius manifolds and moduli spaces for singularities*, volume 151 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2002.
- [19] Nigel Hitchin. Frobenius manifolds. In *Gauge theory and symplectic geometry (Montreal, PQ, 1995)*, volume 488 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 69–112. Kluwer Acad. Publ., Dordrecht, 1997. With notes by David Calderbank.
- [20] H. Iritani. Quantum D-modules and generalized mirror transformations. [math.DG/0411111](https://arxiv.org/abs/math.DG/0411111).
- [21] M. Kapranov and E. Vasserot. Kleinian singularities, derived categories and Hall algebras. *Math. Ann.*, 316(3):565–576, 2000.
- [22] M. Kontsevich and Yu. Manin. Gromov-Witten classes, quantum cohomology, and enumerative geometry. *Comm. Math. Phys.*, 164(3):525–562, 1994.
- [23] A. G. Kouchnirenko. Polyèdres de Newton et nombres de Milnor. *Invent. Math.*, 32(1):1–31, 1976.

- [24] Jun Li and Gang Tian. Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties. *J. Amer. Math. Soc.*, 11(1):119–174, 1998.
- [25] Yuri I. Manin. *Frobenius manifolds, quantum cohomology, and moduli spaces*, volume 47 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1999.
- [26] Paul Seidel. More about vanishing cycles and mutation. In *Symplectic geometry and mirror symmetry (Seoul, 2000)*, pages 429–465. World Sci. Publishing, River Edge, NJ, 2001.
- [27] K. Ueda. Stokes matrices for the quantum cohomologies of Grassmannians. [math.AG/0503355](#).
- [28] Eric Zaslow. Solitons and helices: the search for a math-physics bridge. *Comm. Math. Phys.*, 175(2):337–375, 1996.

Research Institute for Mathematical Sciences, Kyoto University, Oiwake-cho, Kitashirakawa, Sakyo-ku, Kyoto, 606-8502, Japan.
e-mail address : kazushi@kurims.kyoto-u.ac.jp